

ORIGINAL RESEARCH ARTICLE

Evaluation of *Mentha arvensis* L. cultivars for essential oil yield in Southern tropical regions of India

CHANNAYYA HIREMATH^{1*} • YOGENDRA ND¹ • RAMESH VADATTI¹ • K BASKARAN¹
• MANOJ K YADAV¹ • RAM S VERMA² • VELUSAMY SUNDARESAN¹

Article History

Received: December 13th, 2019

Revised: February 11th, 2020

Accepted: March 4th, 2020

Key Words

Cultivars

Essential Oil

Menthol mint

Southern India

ABSTRACT

A field experiment was conducted during 2017-18 at CSIR-CIMAP Research Centre, Bengaluru to study the oil yields of *Mentha arvensis* L. cultivars viz., CIM-Saryu, CIM-Kranti and Kosi for assessing their suitability of cultivation under southern climatic condition. The experiment was laid out in a Randomized Complete Block Design with three treatments, replicated seven times. The three cultivars were hydro distilled for ascertaining their essential oil contents and analysis for quality evaluation. The results revealed that the oil content in Kosi and CIM-Kranti was 1.30%, while that of CIM-Saryu was 1.1 %. Gas Chromatography-Mass Spectrometry (GC-MS) was performed for analysing the menthol content in the essential oils, which showed that there was no significant variation in the menthol content in the three cultivars. The menthol content in Kosi, CIM-Kranti, and CIM-Saryu were estimated to be 77.59%, 78.68 % and 77.86 %, respectively.

© CSIR-Central Institute of Medicinal and Aromatic Plants, Lucknow-226015

INTRODUCTION

Mentha species produce some of the most economically important essential oils throughout the world. Commonly grouped as mints, the genus belongs to the family Lamiaceae (Kumar et al., 2011). *Mentha* comprises a group of 25-30 species (Ali et al., 2002). These plants are of great economic importance since ancient times, because of their therapeutic, culinary and aromatic properties (Dorman et al., 2003). Distillation of aerial parts of these herbs yield an essential oil containing a large number of terpene compounds such as menthol, menthone, isomenthone etc. Angola, Australia, Argentina, Bulgaria, Brazil, France, Hungary, Italy,

Czechoslovakia, Paraguay, India, USA, Thailand and Switzerland are main producers of mint oils (Bahl et al., 2000).

Among these *Mentha* species, *Mentha arvensis* (menthol mint) is an important crop for the plains of North India, providing livelihood to the thousands of farmers across Haryana, Uttar Pradesh, and Punjab. It occupies more than 16000 hectares of cultivation area, and account for nearly 90% of the total menthol mint oil production of the world (Singh et al., 2005; Khanuja, 2007). Under North Indian climatic conditions, *M. arvensis* is mainly cultivated as a spring–summer crop. In recent years, an increasing need for the natural

*Corresponding author; E-mail: channayyahiremath@cimap.res.in

¹CSIR-Central Institute of Medicinal and Aromatic Plants Research Centre, Bangalore, India

²CSIR-Central Institute of Medicinal and Aromatic Plants, CIMAP-Lucknow, India

Doi: <https://doi.org/10.62029/jmaps.v42i2.Hiremath>

menthol of its oil for various industrial requirements has lead to the increased demand for oil production. Mint oil production can be increased through the development of high-yielding varieties, and by increasing the cultivation area beyond the traditional cropping limits with suitable cultivation strategies and packages.

Three most popular menthol mint varieties under commercial cultivation in the North Indian region are Kosi, CIM-Kranti and CIM-Saryu. CIM-Kranti is a cold tolerant genotype, suitable for generating extra income without any additional input and extra land use for cultivation during both winter (100 kg/ha oil yield and 250-300 q/ha suckers) as well as summer (170-210 kg/ha oil yield) seasons. Kosi is an early maturing variety which is tolerant to the leaf spot and powdery-mildew diseases in mint. It has the ability to produce a herb yield of 260 q/ha with an oil yield of 200 kg /ha. CIM-Saryu is one of the oldest and robust varieties, producing a huge biomass and large canopy cover. These three menthol mint varieties viz., Kosi, CIM-Kranti and CIM-Saryu were selected for the present study to evaluate the essential oil content and quality in the non-traditional Southern tropical climatic conditions of Karnataka, India.

MATERIALS AND METHODS

Location of experimental site

Field experiment was conducted at CSIR-CIMAP Research Centre, Bengaluru. The experimental area lies between 13°081' longitude North and 77°571' East at an altitude of 930 meters above mean sea level, which receives an average annual rainfall of about 928 mm.

Experimental Details

Field experiment was laid out in Randomized Complete Block Design during 2017-2018 with three treatments (CIM-Saryu, CIM-Kranti and Kosi), replicated seven times. The varieties were planted in the first week of March, at a spacing of 45×45 cm. The experiment was conducted following the recommended agricultural practices.

Plant sampling for distillation

The crop was harvested at maturity stage for the extraction of essential oil. Harvested fresh herb was hydro-distilled with the help of Clevenger's apparatus for 3 hrs. The oil samples were dried over anhydrous sodium sulphate and stored in refrigerator (4°C) until further analysis.

Essential oil analysis

Gas Chromatography (GC) analysis of the essential oil was performed on a Centurion Scientific Gas Chromatograph (model CS-5800), equipped with FID and a polar (Supelcowax-10) fused silica capillary column (30 m × 0.25 mm internal diameter, film thickness 0.50 µm). Nitrogen was used as the carrier gas at 1.0 ml/min. Oven temperature was programmed from 70°C to 170°C with ramp of 4°C/min, and then programmed to 240°C with ramp of 5°C/min with initial and final hold times of 5 and 15 min, respectively. Injector and detector temperatures were set at 240°C and 250°C, respectively. The samples of 0.1 µL were injected in the split mode (1:70). Identification of the essential oil constituents was carried out on the basis of retention index (RI) determined with reference to homologous series of *n*-alkanes (C₇-C₃₀) and mass spectral data. The relative amounts of individual components were calculated based on the relative % peak areas (FID response), without using correction factor.

Statistical analysis

The data was analyzed using SPAR-2 software (Ahuja et al., 2008) and graphical representation was performed using GraphPad Prism version 5.00 for Windows (GraphPad Software, La Jolla California USA, www.graphpad.com.).

RESULTS AND DISCUSSION


Essential oil content

Distillation of whole herb of the three tested varieties yielded an oil content ranged from 1.10% to 1.30%. CIM-Kranti and Kosi performed better while CIM-Saryu recorded a lower oil content of 1.10% (Fig. 1). Analytical results revealed that there is no significant statistical difference among the

Table 1. Analysis of variances for oil content (%)

SV	df	MSS	F(Cal)	P-value (5%)	F (Critical)	CV (%)
Replication	6	0.009	1.158	0.389	2.996	
Treatments	2	0.075**	9.916	0.003	3.885	8.470
Error	12	0.008				


SV: Sources of variation, df: Degrees of freedom MSS: Mean Sum of squares, CV: Coefficient of variance

**Figure 1:** Mean values of essential oil content (%)

three cultivars of Menthol mint under Bangaluru condition for the oil content (Table 1). However, the varieties CIM-Kranti and Kosi yielded more essential oil than CIM-Saryu.

Essential oil composition

The essential oils obtained upon hydro-distillation of harvested herbs were subjected to GC analysis. Around 20-30 constituents were identified that accounted for 92-95% of the total oil. Menthol, menthone, limonene and isomenthone (Monoterpenoids) were identified as the main constituents in the essential oil of all the three tested varieties. The major volatile constituents of these

**Figure 2:** Chromatogram of a) CIM-Saryu, b) CIM-Kranti and c) Kosi varieties**Table 2. Chemical composition of *Mentha arvensis* essential oil of the three tested varieties under southern tropical conditions**

Compound	CIM-Saryu (%)	CIM-Kranti (%)	Kosi (%)
Limonene	2.2	2.8	3.2
1,8-Cineole	0.1	0.1	0.2
Menthone	4.9	7.3	7.8
Isomenthone	2.8	3.1	3.7
Menthyl acetate	1.9	1.4	1.7
Menthol	77.86	78.68	77.59

oil samples are presented in Table 2. Menthol is the major compound present in all the cultivars (Fig. 2). The highest menthol content recorded is in CIM-Kranti (78.68 %) followed by CIM-Saryu (77.86 %) and Kosi (77.59%). The menthone content in the three cultivars varies between 4.98-7.80 %, being the highest in Kosi (7.80 %) followed by CIM-Kranti (7.6%), and Sarayu (4.5%).

Evaluation of *Mentha arvensis* cultivars under southern tropical regions of India showed that no significant quantitative variation exist among them for their oil content. According to earlier research reports, essential oil yield in aromatic plants are influenced by several agricultural and environmental factors such as nutrient availability, meteorological variations, soil micro-flora, genetic factors, and geographical variables that influences the expression of genes controlling growth and development (Verma et al., 2016; Jamzad et al., 2013; Martin and Chang, 2013). The data pertaining

mid-hills of Uttarakhand (Verma et al., 2010; Padalia et al., 2015), foothill conditions (Singh et al., 2015), western Himalayan region (Padalia et al., 2013). *Mentha arvensis* cv. Shivalik grown in the tropical climate of India shows menthol (53.2-82.3%), menthone (5.2-30.2%), isomenthone (2.1-3.5%) and neomenthol (0.9-2.0%) as major constituents (Rao et al., 2000). The genotype and agronomic conditions, such as harvesting time, plant age and crop density greatly influence the quantitative composition of the essential oils of many aromatic crops (Verma et al., 2010).

It can be concluded from the present study that the quality of the essential oil obtained from three cultivars, viz Kosi, CIM-Kranti and CIM-Saryu in southern tropical region of India are comparable with that obtained in the North Indian plains. Both CIM-Kranti and Kosi are comparatively better suited for the southern climatic conditions, as they yielded a better essential oil content than CIM-Saryu.

Table 3. Comparison of the major chemicals components (% composition) of menthol mint oil in previously reported studies (Akhil and Subhan, 2012)

Constituent	Karnataka	Andhra Pradesh	Tamil Nadu	Madhya Pradesh	Uttar Pradesh	Uttarakhand
3-Octanol	0.10-0.20	0.10	0.90	1.40	-	0.00-0.50
Limonene	2.20-3.20	1.00	-	0.70	-	0.00-0.30
Menthone	4.90-7.80	9.60	15.00	25.10	4.00-11.70	0.30-7.90
Isomenthone	2.80-3.70	4.00	4.60	-	2.00-4.20	0.00-6.10
Menthol	77.50-78.68	73.00	69.50	48.80	72.30-83.80	77.50-89.30
Menthyl acetate	1.40-1.90	4.00	0.90	1.80	2.00-5.90	0.00-0.80
Cultivar	Kosi, CIM-Saryu and CIM-Kranti	Shivalik	Shivalik	-	6 cultivars	6 cultivars
Reference	Present study	Rajeswara Rao (1999a)	Rajeswara Rao et al., (1999b)	Pandey and Chowdhury 2000	Shasany et al., 2010	Singh et al., 2005

to the GC analysis of the essential oil samples (Fig. 2) obtained by us also illustrate that menthol is the major chemical constituent, followed by menthone. Even though statistically there is not much qualitative difference in the major essential constituents and oil content in all three varieties, considerable variations in the quantitative make-up of the essential oil constituents were observed. Similar results are also reported by many researchers (Table 3). Menthol, menthone, isomenthone, limonene and methyl acetate are the major constituents in *Mentha arvensis* from the

ACKNOWLEDGEMENT

The authors are grateful to the Director, CIMAP, Lucknow (UP), India, for continuous encouragement and providing necessary facilities.

REFERENCES

- Ahuja S, Malhotra PK, Bhatia VK, Rajender Parsad 2008. Statistical Package for Agricultural Research (SPAR 2.0). *J Ind Soc Agril Statist* **62**: 65-74.

- Akhil B, Subhan CN, 2012. *Natural Essential Oils Fragrances and Flavours*, Aavishkar Publishers, Distributors Jaipur 3A2 003 (Raj.). India.
- Ali MS, Saleem M, Ahmad W, Parvez M, Yamdagni RA. 2002. Chlorinated monoterpene ketone, acylated_-sitosterol glycosides and a flavanone glycoside from *Mentha longifolia* (Lamiaceae). *Phytochem* **59**: 889–895.
- Bahl JR, Bansal RP, Garg SN, Naqvi AA, Luthra R, Kukreja AK, Kumar S. 2000. Qualitative evaluation of the essential oils of the prevalent cultivars of commercial mint species *Mentha arvensis*, *M. spicata*, *M. piperita*, *M. cardiaca*, *M. citrata* and *M. Viridis* cultivated in indo-gangetic plains. *J Med Arom Pl Sci* **22**: 787–797.
- Dorman HD, Kosar M, Kahlos K, Holm Y, Hiltunen R. 2003. Antioxidant properties and composition of aqueous extracts from *Mentha* species, hybrids, varieties, and cultivars. *J Agric Food Chem* **51**: 4563–4569.
- Jamzad MZ, Jamzad F, Mokhber S, Ziareh, Yari M. 2013. Variation in essential oil composition of *Mentha longifolia* var. *Chlorodictya* Rech. f. and *Ziziphora clinopodioides* Lam. growing in different habitats. *J Med Pl Res* **7**: 1618–1623.
- Khanuja SPS. 2007. Aroma vision 2020: Technology path of cultivating today for a fragrant tomorrow. In: Proc. National Convention & Seminar on Business Enabling of Aromatic Plants & Products, Dehradun, India. November 21–22.
- Kumar P, Mishra S, Malik A, Satya S. 2011. Insecticidal properties of *Mentha* species: A review. *Ind Crop Prod* **34**: 802–817.
- Martin P, Chang X. 2013. Production potential and crop agronomy of sweet gale (*Myrica gale* L.) in the north of Scotland. *Ind Crop Prod* **46**: 39–49.
- Padalia RC, Verma RS, Chauhan A, Sundaresan V, Chanotiya CS. 2013. Essential oil composition of sixteen elite cultivars of *Mentha* from western Himalayan region, India Maejo Int. *J Sci Technol* **7**: 83–93.
- Padalia RC, Verma RS, Chauhan A, Goswami P. 2015. Quality evaluation of elite mint cultivars in two cropping seasons in subtropical conditions of Uttarakhand, India, *J Essent Oil Res* **27**: 45–51.
- Pandey AK, Chowdhury AR. 2000. GC-MS studies of Japanese mint (*Mentha arvensis* L.) oil. *J Med Arom Pl Sci* **22**: 468–469.
- Rao BRR, Bhattacharya AK, Singh K, Kaul PN, Mallavarapu GR. 1999. Comparative composition of cornmint oils produced in north and south India. *J Ess Oil Res* **11**: 54–56.
- Rao BRR, Kaul PN, Bhattacharya AK, Naqui AA. 1999a. Cultivation of cornmint (*Mentha arvensis* L. f. *piperascens* Malinv. Ex Holmes) in Tamil Nadu. Yield quality of the essential oil. *J Essent Oil Bear Plants* **2**: 127–131.
- Rao BRR, Kaul PN, Mallavarapu GR, Ramesh S. 2000. Comparative composition of whole herb, flowers, leaves and stem oils of cornmint (*Mentha arvensis* L.f. *piperascens* Malinvaud ex Holmes), *J Essent Oil Res* **12**: 357–359.
- Shasany AK, Gupta S, Gupta MK, Naqvi AA, Bala JR, Khanuja SPS. 2010. Assessment of menthol mint collection for genetic variability and monoterpene biosynthetic potential. *Favour Fragr J* **25**: 41–47.
- Singh AK, Raina VK, Naqvi AA, Patra NK, Kumar B, Ram P, Khanuja SPS. 2005. Essential oil composition and chemo-arrays of menthol mint (*Mentha arvensis* L. f. *piperascens* Malinvaud ex. Holmes) cultivars, *Flav Fragr J* **20**: 302–305.
- Verma RS, Rahman L, Verma RK, Chauhan A, Yadav AK, Singh A. 2010. Essential oil composition of menthol mint (*Mentha arvensis*) and peppermint (*Mentha piperita*) cultivars at different stages of plant growth from Kumaon region of Western Himalaya. *Open Acc J Med Arom Pl* **1**: 13–18.